

Summary of recent (heavy ion) results from LHC



Tatsuya Chujo

2013 RHIC & AGS Annual Users' Meeting

June 27, 2013, BNL, USA



筑波大学
University of Tsukuba

Outline



- ◆ Introduction
- ◆ First three years of LHC heavy ion runs
- ◆ Highlights from p-Pb results (2013)
- ◆ Summary



13 yeas ago..., at BNL

June 12, 2000 @ PHENIX

First collisions at RHIC at $\sqrt{s_{NN}} = 56 \text{ GeV Au+Au}$






2009.11.23 at the CERN-ALICE control room



ALICE

**First proton-proton collisions
 $p+p \sqrt{s} = 900 \text{ GeV}$**

First proton-proton collisions p+p \sqrt{s} = 900 GeV in ALICE (2009.11.23)





Recognized by European Physical Society

Particles and Fields


First proton–proton collisions at the LHC as observed with the ALICE detector: measurement of the charged-particle pseudorapidity density at $\sqrt{s} = 900$ GeV

First ALICE
publication

submitted to EPJC 28 Nov 2009

Eur. Phys. J. C (2010) 65: 111–125
DOI 10.1140/epjc/s10052-009-1227-4

arXiv:0911.5430v2




From the ALICE Collaboration: First proton–proton collisions at the LHC as observed with the ALICE detector: measurement of the charged particle pseudorapidity density at $\sqrt{s} = 900$ GeV.

Società Italiana
di Fisica

Springer

LHC First Physics

30 March 2010






7 TeV collision events seen today by the LHC's four major experiments (clockwise from top-left: ALICE, ATLAS, CMS, LHCb). [More LHC First Physics images »](#)

LHC research programme gets underway

Geneva, 30 March 2010. Beams collided at 7 TeV in the LHC at 13:06 CEST, marking the start of the LHC research programme. Particle physicists around the world are looking forward to a potentially rich harvest of new physics as the LHC begins its first long run at an energy three and a half times higher than previously achieved at a particle accelerator. [Read more...](#)






Mar. 30, 2010

First p-p collisions
at $\sqrt{s} = 7 \text{ TeV}$



Nov. 8, 2010


First Pb-Pb collisions at LHC,
the opening new era of
heavy ion program at LHC





LHC run history

- **2009:** Commissioning and first data p-p (900 GeV)
- **2010:** First p-p run (7 TeV) and **first Pb-Pb run (2.76 TeV)**
- **2011:** Long p-p (7 TeV) and **one month Pb-Pb (2.76 TeV)** = x10 luminosity than that in 2010. first p-p (2.76 TeV).
- **2012:** Long p-p (8 TeV), one day p-Pb (5.02 TeV) pilot run
- **2013:** 1.5 month p-Pb and Pb-p run (5.02 TeV), (32 nb⁻¹ in ALICE)
- **2013.02 - 2014 winter:** LHC Long Shutdown I (LSI) ← We are here now



First three years of LHC heavy ion runs



Initial temperature

Direct photon p_T spectra at LHC


- Observed an excess over p-p baseline at low p_T (< 2 GeV/c).


$T = 304 \pm 51$ MeV

- ~30% higher than RHIC



RHIC (200 GeV Au+Au):
 $T = 221 \pm 19 \pm 19$ MeV

PHENIX, PRL 104, 132301 (2010)






Dissociation temperature



CMS, PRL 109
(2012) 222301




Melting excited Υ states

- Suppression of ground state $Y(1s)$, and excited states $Y(2S)$ and $Y(3S)$.
- Consistent with **the sequential melting scenario**, $Y(3S) > Y(2S) > Y(1S)$.




ALICE

Freeze-out T_{kin} and $\langle\beta_T\rangle$



ALICE, PRL, 109 252301 (2012)
ALICE, arXiv:1303.0737

Significant changes in slope compared to RHIC, especially for protons.



ALI-PUB-47066

Blast-wave fits


- $T_{\text{kin}} = 95 \pm 10 \text{ MeV}$
→ comparable with RHIC
- $\langle\beta_T\rangle = 0.65 \pm 0.02$
→ 10% higher than RHIC



ALICE

T_{ch} and μ_b

- Measured dN/dy of PID hadrons at mid-rapidity at LHC.
- Data: feed-down corrected.






ALICE

T_{ch} and μ_b

- Measured dN/dy of PID hadrons at mid-rapidity at LHC.
 - Data: feed-down corrected.
 - Thermal statistical model with $T_{ch} = 164$ MeV, $\mu_b = 1$ MeV
- ➔ does not reproduce the data well, especially p, Λ, ϕ, K^*






ALICE

T_{ch} and μ_b


- Measured dN/dy of PID hadrons at mid-rapidity at LHC.
- Data: feed-down corrected.
- Thermal statistical model with $T_{ch} = 164 \text{ MeV}$, $\mu_b = 1 \text{ MeV}$
 - ➔ does not reproduce the data well, especially p, Λ, ϕ, K^*
- $T_{ch} = 152 \text{ MeV}$, $\mu_b = 1 \text{ MeV}$ w/o ϕ, K^* , improving the fit, but multi-strangeness (Ξ, Ω) does not get right.



Indicating the importance of **re-scattering at hadronic phase?**



Energy density



Multiplicity density:

2.1x Central Au-Au at 200 GeV

Energy density:



~ 3 x RHIC (larger $\langle m_T \rangle$)

$$\epsilon\tau \approx 16 \text{ GeV}/(\text{fm}^2 c)$$

B. Mueller et al., Ann.Rev.Nucl.Part.Sci.62 (2012) 361
ALICE: PRL 106 (2011) 032301
CMS: JHEP 1108 (2011) 141
ATLAS: PLB 710 (2012) 363

Freeze-out volume and lifetime



- Freeze-out volume: **$300 \text{ fm}^3 \sim 2 \times \text{RHIC}$** .
- Lifetime: **$10 \text{ fm/c} \sim 40\% \text{ longer than that at RHIC}$** .




$$\tau_f = R_{\text{long}} \sqrt{m_T/T}$$




Elliptic flow v_2 at LHC




- 30 % increase compared to RHIC data, due to $\langle p_T \rangle$ increase.
- p_T dependence holds from RHIC to LHC.
- **Suggesting similar η/s at LHC as RHIC produced.**




v_n and initial conditions, η/s



- v_n measurements by two particle correlations, $|\Delta\eta| > 2$, at very central collisions (0-0.2%).



v_n and initial conditions, η/s



← Glauber with different η/s

- Power spectrum of v_n .
- disentangle initial condition and η/s .


← Glauber and CGS with η/s
VISH2+1 Hydro

CMS-HIN-12-011



Further constraint on η/s ; E-by-E v_n

ATLAS-CONF-2012-114




Theory: arXiv:nucl-ex/0701025, Phys. Rev. C 74, 044905 (2006)

- Direct measurements of v_2, v_3, v_4 (only v_2 shown).
- Model comparison:
 - both work in 0-1%
 - **MC-KLN (CGC)** works in 5-10%
 - **Glauber** works in 40-45%
- Additional constraints by event-plane correlations (ATLAS-CONF-2012-049).




Quark number scaling of v_2

LHC



ALICE-PREL-12345




RHIC



PHENIX, PRL 98, 162301 (2007)


- K_{ET}/n_q scaling at LHC does not work well like those at RHIC.
- Affected by a strong radial flow for protons (hadronic re-scatterings)?

Jets in LHC heavy ion collisions






Energy loss, single hadron R_{AA}



**Baryon
enhancement
($p_T < 8$ GeV/c)**

**recombination
+ radial flow**

R_{AA} for π, K, p





- EM probes: $R_{AA} \sim 1$.
- Hadrons: strong suppression $R_{AA} \sim 0.1$ at ~ 7 GeV/c, and a rise at higher p_T ($R_{AA} \sim 0.5$).
- Baryon enhancement is observed at intermediate p_T .




Energy loss, jet R_{AA}

ALICE (30-120 GeV)
Full jets (0-10%)



CMS (30-120 GeV)
Full (q/g) jets (0-5%)





- a wide kinematic reach for jets at LHC.
- Jet $R_{AA} \sim 0.5$ above 100 GeV/c, consistent with hadron's R_{AA} .



Energy loss, jet R_{AA}

ALICE (30-120 GeV)
Full jets (0-10%)

CMS (30-120 GeV)
Full (q/g) jets (0-5%)




- Jet R_{AA} from $p_T = 30 - 300$ GeV, consistent with hadron's R_{AA} .



Di-jet energy imbalance

CMS, PRC 84, 024906 (2011)




Large energy imbalance is observed in central Pb-Pb.

$$A_J = \frac{p_{T,1} - p_{T,2}}{p_{T,1} + p_{T,2}}$$

$p_{T,1}$: leading jet
 $p_{T,2}$: sub-leading jet

Large A_J : low momentum particle (< 4 GeV/c) emitted at large angle on away side.

ATLAS, PRL, 105 (2010) 252303







γ -jet: jet tomography

CMS, Phys. Lett. B 718 (2013) 773

$$\langle x_{J\gamma} \rangle = p_T^{jet} / p_T^\gamma$$

$R_{J\gamma}$: fraction of photons with jet partner





- γ as a calibrated probe of jet energy.
- significant change in $R_{J\gamma}$, $\langle x_{J\gamma} \rangle$ compared to PYTHIA and pp.




ALICE

Heavy quark R_{AA}



- D mesons are also strongly suppressed.
- **A hit of $R_{AA}^D > R_{AA}^{\pi}$** (not yet conclusive).

Charm vs. Bottom




* Note: not same kinematic range.

- R_{AA} for charmed meson (D) vs. bottom meson (J/ψ from B decay).
- First indication of a mass dependence of R_{AA} .
- $R_{AA}^B > R_{AA}^D$

J/ψ (color screening vs. regeneration)


mid-rapidity R_{AA} for J/ψ



- J/ψ measured at mid-rapidity $|y| < 0.9$, by e^+e^- at LHC.
- Compared to RHIC mid-rapidity data.
- Significant larger R_{AA} than those at RHIC.

J/ψ (color screening vs. regeneration)

forward-rapidity R_{AA} for J/ψ




- J/ψ measured at forward-rapidity $2.5 < y < 4$, by $\mu^+\mu^-$ at LHC.
- Compared to RHIC forward data.
- Significant larger R_{AA} than those at RHIC.
- Suppression is stronger than that at mid-rap.





J/ψ (color screening vs. regeneration)


Low p_T : R_{AA} at forward y , $J/\psi \rightarrow \mu^+\mu^-$



High p_T : R_{AA} at forward y , $J/\psi \rightarrow \mu^+\mu^-$




vs.



- J/ψ R_{AA} is enhanced at low p_T .
- Compatible with models including regeneration.

Charmonia flow (Inclusive J/ψ v_2)

ALICE: arXiv:1303.5880



- J/ψ produced via regeneration of thermal de-confined c-quarks should show **a non zero v_2** .
- **First hint of non-zero v_2** .
- Consistent with the transport model with regeneration.

Highlights from p-Pb results (2013)







ALICE

$dN/d\eta$, R_{AA} in p-Pb

ALICE, PRL 110 (2013) 032301



ALICE, PRL 110 (2013) 082302




- $dN/d\eta$: most models reproduce data <20%. CGC: steeper rise on shape.
- $R_{AA} \sim 1$ in pPb: suppression in Pb-Pb central is a final state effect.



Di-Hadron Correlations in p-p & p-Pb

p-p ($N \geq 110$)


CMS $N \geq 110$, $1.0 \text{ GeV}/c < p_T < 3.0 \text{ GeV}/c$




p-Pb ($N \geq 110$)

CMS pPb $\sqrt{s_{NN}} = 5.02 \text{ TeV}$, $N_{\text{trk}}^{\text{offline}} \geq 110$

$1 < p_T < 3 \text{ GeV}/c$



$\Sigma E_T^{\text{Pb}} < 20 \text{ GeV}$




p-Pb ($\Sigma E_T^{\text{Pb}} < 20 \text{ GeV}$)

ATLAS p+Pb $\sqrt{s_{NN}} = 5.02 \text{ TeV}$

$\int L \approx 1 \mu\text{b}^{-1}$, $0.5 < p_T^{a,b} < 4 \text{ GeV}$

(a)

$\Sigma E_T^{\text{Pb}} > 80 \text{ GeV}$



p-Pb ($\Sigma E_T^{\text{Pb}} > 80 \text{ GeV}$)

- First observation of **ridge structure in high multiplicity p-p** (CMS).
- Also confirmed in **p-Pb high multiplicity events**.
- Always side ridge structure is observed in high multiplicity p-Pb.


CMS, JHEP 1009 (2010) 91
CMS, PLB 718 (2012) 795
ATLAS, PRL 110, 182302 (2013)




ALICE

Double ridge structure in p-Pb

ALICE, PLB 719 (2013) 29




0-20%




60-100%



=



- Extract double ridge structure by subtracting p-p jet like distribution in p-Pb (60-100%) from central p-Pb (0-20%).
- Confirmed that near and away side ridges are almost same structure.
- **Strong correlation between near and away side yields, suggesting the same origin.**



J/ψ R_{pPb}





Comparison between prompt and inclusive J/ψ:

- Measurements are consistent within uncertainties, although prompt is $\sim 30\%$ lower overall.
- Provides further constraints on CGC model.

Summary

- Hottest, largest, longer lived QGP is produced at LHC heavy ion collisions.
- There are similarities, differences compared to RHIC, and newly discovered properties on QGP at LHC.
- We enter an era of determination of QGP properties by jets, photons, c/b quarks, quarkonium with bulk particles.
- **Future**
 - Run with full energy $\sqrt{s_{NN}} = 5.5$ TeV Pb-Pb in 2015-2017 (Run-2), with upgraded detectors (LSI, ALICE).
 - Preparing the detector upgrade for higher luminosity LHC run during LS2 (2018) for Run-3 (2019-2022).



**Thank you for your
attentions!**

